

Fluorescent entity comprising a fluorophore covalently attached to at least one oligonucleotide and comprising at least one functional group, and uses thereof

5 The invention relates to a fluorescent entity comprising a fluorophore, covalently attached to one or more oligonucleotides or oligonucleotide analogs and comprising at least one functional group, introduced or generated on the fluorophore or one of the oligonucleotides or
10 oligonucleotide analogs.

Many families of organic molecules are used as fluorescent labels for biomolecules in a number of applications, in particular for diagnostic methods which
15 make it possible to follow or to quantify these biomolecules.

Mention may in particular be made of rhodamines, cyanins, squaraines or bodipy dyes. Most of these molecules have a
20 high molar extinction coefficient (often greater than $100\,000\text{ M}^{-1}\text{ cm}^{-1}$) and a quantum yield of fluorescence generally greater than 20%.

However, due to their often very hydrophobic nature,
25 these organic molecules have a tendency to form aggregates at the surface of biomolecules, in particular proteins, when the desire is to have several fluorescent labels per protein (U. Schobel et al., Bioconjugate chem., 1999, 10, 1107-1114). Since these aggregates are
30 virtually nonfluorescent, the mean quantum yield of the fluorescent molecules present at the surface of the proteins is then significantly lower than that of the native molecules.

35 To overcome this problem, an attempt has been made, in the literature, to increase the hydrophilic nature of the

fluorescent molecules, in particular by adding sulfonate groups (US 5,268,486). Mention may also be made of the addition of sugars or of carbohydrate residues to the structure of the fluorescent molecules (WO 98/49176).

5 However, the addition of these units limits the aggregation phenomenon without suppressing it.

Other approaches have also been used to avoid the decrease in quantum yield of fluorescence after labeling
10 of proteins. Application WO 98/26287 describes a method using cyclodextrins to encapsulate fluorescent molecules at the surface of proteins and to limit their aggregation. However, this technique does not function satisfactorily with all types of molecules.

15

A technique has recently been described which makes it possible to obtain proteins having, at their surface, only a single fluorescent molecule and makes it possible to thus be able to maintain a high quantum yield by
20 avoiding the aggregation phenomenon (Winckler et al., Specific labeling of proteins using reactive affinity tag-dye systems, SBS Conference, Vancouver, 2000). This system is limiting since it does not make it possible to have several fluorescent molecules per protein.

25

The problem to be solved therefore consists in providing a label which can be attached to a biomolecule, the photophysical properties of which are not modified, in particular by aggregation phenomena, when several of
30 these labels are simultaneously attached to a biomolecule, in particular a protein.

According to the invention, this problem can be solved by attaching the fluorophore to one or more oligo-
35 nucleotide(s) or oligonucleotide analog(s), the compound thus formed also comprising one or more reactional

group(s) allowing it to be attached to a carrier molecule.

According to a first aspect, the invention therefore
5 relates to a fluorescent entity comprising a fluorophore,
with the exception of a rare earth metal cryptate,
covalently attached to one or more oligonucleotide(s) or
oligonucleotide analog(s), characterized in that it
10 comprises at least one functional group, introduced or
generated on the fluorophore or on one of the oligo-
nucleotides or oligonucleotide analogs.

The fluorophore of the entity according to the invention
preferably comprises one or more aromatic rings and has a
15 high molecular extinction coefficient, greater than
20 000, preferably greater than 50 000.

Said fluorophore is preferably chosen from rhodamines,
cyanins, squaraines, bodipys, fluoresceines and their
20 derivatives.

The term "oligonucleotide" denotes equally oligodeoxy-
ribonucleotides (DNA fragment) or oligoribonucleotides
(RNA fragment).

25 The term "oligonucleotide or oligonucleotide analog" is
intended to mean, in the present description:

- 30 - either a series of ribonucleotide or deoxyribo-
nucleotide units attached to one another via bonds
of the phosphodiester type;
- or a series of ribonucleotide or deoxyribonucleotide
units or of nucleotide analog units modified on the
35 sugar or on the base and attached to one another by
natural internucleotide bonds of the phosphodiester

type, some of the internucleotide bonds being optionally replaced with phosphonate, phosphoramidate or phosphorothioate bonds. These various oligonucleotide families are described in Goodchild,
5 *Bioconjugate Chemistry*, 1(3), May/June 1990, 77-99;

- or a series comprising both ribonucleotide or deoxyribonucleotide units attached to one another by bonds of the phosphodiester type and nucleoside
10 analog units attached to one another by amide bonds, commonly called "PNAs" ("peptide nucleic acids"), as described in M. Egholm et al., J. Am. Chem. Soc., 1992, 114, 1895-1897; such compounds are, for example, described in R. Vinayak et al., Nucleoside
15 & Nucleotide, 1997, 16 (7-9), 1653-1656;

- or a series of ribonucleotide or deoxyribonucleotide units in which some of the nucleosides or of the internucleotide bonds have been modified compared to
20 a natural oligoribonucleotide or oligodeoxyribonucleotide formed from the common nucleosides (adenosine, deoxyadenosine, cytidine, deoxycytidine, guanosine, deoxyguanosine, uridine or thymidine) attached by phosphate bridges, for example a
25 phosphorothioate oligonucleotide (OPT) in which all or part of the phosphate bridges have been replaced with thiophosphate bridges (P.J. Romaniuk, F. Eckstein (1982) J. Biol. Chem. 257, 7684).

30 The use of each one of these types of oligonucleotide constitutes an advantageous aspect of the invention.

The term nucleotide or nucleoside "analog" is intended to mean a nucleotide/nucleoside comprising at least one
35 modification relating to the sugar or the nucleobase or a

combination of these modifications. By way of example, mention may be made of the following modifications:

I. Modifications concerning the sugar (nucleotide or
5 nucleoside analogs):

1°) The sugar component can be modified in that the
configuration of the hydroxyls (free or involved in
a phosphate bridge) is different from the natural
10 configuration (which is, respectively, β -D-erythro
in the DNA series and β -D-ribo in the RNA series),
as in the analogs having the backbone β -D-arabino-
pentofuranoside or β -D-xylo-pentofuranoside, for
example...

15 2°) The structure can be modified in that the
internucleotide bonds are of the 2'→5' type, such
as in the case of β -D-ribo-pentofuranoside-2'-
phosphate or 3'-deoxy- β -D-erythro-pentofuranoside-
20 2'-phosphate derivatives.

Nucleotides exist in which the structure includes
the preceding two modifications, such as β -D-xylo-
pentofuranoside-2'-phosphate.

25 3°) The structure can differ from the natural model in
that the 4' carbon has the opposite configuration,
this is the case α -L-threo-pentofuranoside-3'-
phosphate. The difference may relate to the
configuration of the carbon in the 1'-position
30 (anomeric position), this is the case of α -D-
erythro-pentofuranoside-3'-phosphate. Nucleotides/
nucleosides exist in which the structure includes
the preceding two modifications, such as β -L-threo-
35 pentofuranoside-3'-phosphate.

4°) The structure can differ from the natural model in that the oxygen in the 4'-position is replaced with a carbon (carbocyclic analog) or with a sulfur, such as 4'-thio- β -D-erythro-pentafuranoside-3'-phosphate.

5

5°) The structure can differ from the natural model in that one of the hydroxyls of the sugar is alkylated, for example in the backbone 2'-O-alkyl- β -D-ribo-pentafuranoside-3'-phosphate, the alkyl group possibly being, for example, the methyl or allyl group.

10

6°) The structure can differ from the natural model in that only the sugar component is conserved, such as in 1,2-dideoxy-D-erythro-pentafuranose-3-phosphate, or in that the sugar is replaced with a polyol such as propanediol.

15

II. Modifications concerning the nucleobase (nucleotide analogs):

20

1°) The nucleobase can be modified in that the substituents of the natural bases are modified, such as in 2,6-diaminopurine, hypoxanthine, 4-thiothymine, 4-thiouracil or 5-ethynyluracil.

25

2°) The positions of the substituents can be switched compared to the natural bases, such as in isoguanosine or isocytosine.

30

3°) A nitrogen atom of the nucleobase can be replaced with a carbon, as in the 7-deazaguanosine or 7-deazaadenine.

III. Modifications concerning the internucleotide bond:

Moreover, as mentioned above, the bonds between the sugar units or their analogs can also be modified, for example by replacing one or more of the oxygen atoms of the natural phosphodiester bond with a carbon (phosphonate series), a nitrogen (phosphoramidate series) or a sulfur (phosphorothioates).

- 10 The internucleotide bonds can also be replaced with amide bonds, as in oligonucleotide analogs of the "PNA" type.

According to a preferred aspect, the oligonucleotide consists of a series comprising both ribonucleotide or deoxyribonucleotide units attached to one another by bonds of the phosphodiester type and nucleoside analog units attached to one another by amide bonds.

20 In particular, in this case, said oligonucleotide can comprise at least five internucleotide bonds of the phosphodiester type at the end intended to be attached to the fluorophore.

Preferably, said oligonucleotide or oligonucleotide analog comprises from 5 to 60 nucleotide units, in particular from 5 to 20, preferably 5 to 15 nucleotide units.










30 The fluorescent entity according to the invention should comprise at least one functional group which allows it to be coupled with a carrier molecule.

Advantageously, the functional group is an amine function of a nucleotide unit of the oligonucleotide or of the oligonucleotide analog, or results from the reaction of a free amine function of a nucleotide unit of the

oligonucleotide or of the oligonucleotide analog using a homobifunctional or heterobifunctional reagent which makes it possible to introduce a functional group chosen from the groups: activated ester of a carboxylic acid, 5 carboxylic acid, isothiocyanate, aldehyde, carbonyl, sulfonyl halide, alkyl halide, azide, hydrazide, dichlorotriazine, anhydride, haloacetamide, maleimide and sulfhydryl. The homobifunctional and heterobifunctional reagents and also their use are described in 10 "Bioconjugation" (chapters 5.3 to 5.6, M. Aslam & A. Dent, Macmillan, London, 1998).

Said functional group can, for example, result from the reaction of a free amine function of a nucleotide unit of 15 the oligonucleotide or of the oligonucleotide analog, with an N-hydroxysuccinimidyl ester.




According to a preferred aspect, the functional group is chosen from the groups: maleimide, carboxylic acid, 20 haloacetamide, alkyl halide, azido, hydrazido, aldehyde, ketone, amino, sulfhydryl, isothiocyanate, isocyanate, monochlorotriazine, dichlorotriazine, aziridine, sulfonyl halide, acid halide, hydroxysuccinimide ester, hydroxy-sulfosuccinimide ester, imido ester, hydrazide, azido- 25 nitrophenyl, azidophenyl, azide, 3-(2-pyridyldithio)-propionamide and glyoxal, and more particularly the groups of formula:



where n ranges from 0 to 8 and p is equal to 0 or 1, and Ar is a 5- or 6-membered heterocycle comprising 1 to 3 hetero atoms, optionally substituted with a halogen atom.


- 5 According to an advantageous aspect, the functional group(s) is (are) attached to the fluorophore and/or to the oligonucleotide by a spacer arm consisting of a divalent organic radical, chosen from linear or branched C₁-C₂₀ alkylene groups optionally containing one or more
10 double bonds or triple bonds and/or optionally containing one or more hetero atoms, such as oxygen, nitrogen, sulfur, phosphorus, or one or more carbamoyl or carboxamido group(s); C₅-C₈ cycloalkylene groups and C₆-C₁₄ arylene groups, said alkylene, cycloalkylene or
15 arylene groups being optionally substituted with alkyl, aryl or sulfonate groups.

In particular, the spacer arm is chosen from the groups:



in which n_1 and n_2 are between 2 and 6.

- 5 According to a preferred aspect, the invention relates to a fluorescent entity of formula (I)





(I)


in which:

- A represents a group chosen from:

5










$r = 2 \text{ or } 3$



$r = 2 \text{ or } 3$

- the dashed lines each represent the carbon atoms required to form 1 to 3 fused rings, the groups R_3 being attached to these rings;
- X and Y each represent N, C=O, O, S or $C(CH_3)_2$
- m has a value 1, 2, 3 or 4;
- q has a value 1, 2 or 3;
- $(R_3)_q$ represents q groups R_3 , which may be identical or different;
- the groups R_1 , R_2 and R_3 are identical or different and are chosen from hydrogen; a group $-(CH_2)_s-Z$ in which s ranges from 0 to 4 and Z represents a group CH_3 , SO_3H , OH or $N^+R_1R_2R_3$ in which R_1 , R_2 and R_3 are as defined above; a functional group as defined above; and an oligonucleotide or oligonucleotide analog optionally comprising a functional group as defined above;

- R_4 is chosen from: H; OH; CH_3 ; Cl and the groups of formula:




5

the substituents R_4 in the allylic position possibly forming, with the polyethylenic chain, 1 to 3 fused rings containing from 4 to 14 atoms, which may or may not be saturated, said rings possibly containing one or more atoms of O, N and S, and possibly being optionally substituted with an oxo group.


10

Preferred fluorescent entities according to the invention correspond to formulae (II) and (III)

15




or




5,

in which the dashed lines, R_1 , R_2 , R_3 , R_4 , X , m and q are as defined above for formula (I).


- 10 Advantageously, the invention relates to fluorescent entities of formula (IV), (V), (VI) or (VII)




(IV)



(V)



(VI)




(VII)

in which R_1 , R_2 , R_3 and R_5 are identical or different and
 5 are chosen from hydrogen; a group $-(CH_2)_s-Z$ in which s
 ranges from 0 to 4 and Z represents a group CH_3 , SO_3H , OH
 or $N^+R_1R_2R_3$ in which R_1 , R_2 and R_3 are as defined above; a
 functional group or an oligonucleotide or oligonucleotide
 analog as defined above.

10


According to another preferred aspect, the invention
 relates to a fluorescent entity of formula (VIII)



(VIII)

in which the substituents R_6 to R_{12} are chosen from:
hydrogen; a halogen; an alkyl; a cycloalkyl; aryl;
arylalkyl; acyl; sulfo; a functional group or an
5 oligonucleotide or oligonucleotide analog as defined
above.

Another fluorescent entity according to the invention
corresponds to formula (IX)




(IX)

10

in which R_1 , R_2 , R_3 , R_4 , X , Y , m and q are as defined
above.

15 Entities of formula (IX) which are particularly preferred
are those in which X and Y represent a group $C(CH_3)_2$, and
also those in which

20 - R_1 and R_2 represent an alkyl comprising from 1 to 4
carbon atoms or a group of formula below, at least
one of the groups R_1 and R_2 representing a group of
formula below:










- R_4 represents hydrogen
- $q = 1, m = 2$

5 R_3 represents hydrogen; a group $-(CH_2)_s-Z$ in which s ranges from 0 to 4 and Z represents a group CH_3 , SO_3H , OH or $N^+R_1R_2R_3$ in which R_1 , R_2 and R_3 are as defined above; a functional group or an oligonucleotide or oligonucleotide analog as defined above;

10

- R_4 is chosen from: H ; OH ; CH_3 ; Cl and the groups of formula:






15

the substituents R_4 in the allylic position possibly forming, with the polyethylenic chain, 1 to 3 fused rings containing from 4 to 14 atoms, which may or may not be saturated, said rings possibly containing one or more

atoms of O, N and S, and possibly being optionally substituted with an oxo group.

5 Advantageously, the fluorescent entity according to the invention comprises a fluorophore which is covalently attached to the oligonucleotide, either directly or via a spacer arm.

10 This spacer arm may, for example, consist of a divalent organic radical chosen from linear or branched C₁-C₂₀ alkylene groups optionally containing one or more double bonds or triple bonds and/or optionally containing one or more hetero atoms, such as oxygen, nitrogen, sulfur, phosphorus, or one or more carbamoyl or carboxamido
15 group(s); C₅-C₈ cycloalkylene groups and C₆-C₁₄ arylene groups, said alkylene, cycloalkylene or arylene groups optionally being substituted with alkyl, aryl or sulfonate groups, or chosen from the groups:



in which n_1 and n_2 are between 2 and 6.

- 5 According to a subsequent aspect, the invention also relates to the fluorescent conjugates consisting of an entity as defined above covalently attached to a carrier molecule.
- 10 Advantageous conjugates are those in which the final molar ratio, defined as the number of moles of fluorescent entities per carrier molecule, is greater than 0 and less than 100, preferably less than 20.

The carrier molecule is, for example, an antibody, an antigen, an intracellular messenger, an intercellular messenger, a protein, a peptide, a hapten, a lectin, biotin, avidin, streptavidin, a toxin, a carbohydrate, an oligosaccharide, a polysaccharide, a nucleic acid, a hormone, a vitamin, a medicinal product, a polymer, a polymeric particle, glass, a particle of glass or a surface made of glass or of a polymer.

10 The use of the fluorescent entities according to the invention makes it possible to produce conjugates exhibiting virtually zero aggregation of the fluorophore. Consequently, the quantum yield of the fluorophore can be almost completely conserved after attachment to carrier
15 molecules, even when the final molar yield (number of moles of fluorescent entities per carrier molecule) increases.

This makes these conjugates very advantageous for use in
20 a fluorescent system using nonradiative energy transfer (of the HTRF type). They are also of great advantage in more conventional techniques of detection by fluorescence, where the number of fluorophores per carrier molecule, the quantum yield and the molar
25 extinction coefficient of the fluorophore are predominant criteria for the sensitivity of these systems.

The invention therefore also relates to the use of a fluorescent entity or of a fluorescent conjugate as
30 defined above, as fluorescent tracer(s), for example for detecting and/or determining, by fluorescence, an analyte in a medium liable to contain it or for determining an interaction between biomolecules; or for determining a biological activity such as: an enzyme activity, the
35 activation of a membrane-bound receptor, the transcription of a gene, a membrane transport or a

variation in membrane polarization, in particular in a method for screening medicinal products.

5 The fluorescent conjugates according to the invention can be used as acceptor fluorescent compounds in the presence of donor fluorescent compounds or as donor fluorescent compounds in the presence of acceptor fluorescent compounds, in particular in fluorescence microscopy, in flow cytometry, in fluorescence polarization or in
10 fluorescence correlation.

They can also be advantageously used as a contrast agent for optical imaging in vivo.

15 A subject of the invention is also a method for decreasing the phenomenon of aggregation at the surface of a carrier molecule attached to a fluorophore, characterized in that a fluorescent entity as defined above is used in place of said fluorophore.

20 Finally, a subject of the invention is a method for increasing the quantum yield of a fluorophore attached to a carrier molecule, characterized in that a fluorescent entity as defined above is used as a fluorophore.

25 The fluorescent entities according to the invention can be prepared as described below, by coupling "functionalized" oligonucleotides with a fluorophore.

30 In the present description, the term "functionalized oligonucleotide" is intended to mean an oligonucleotide comprising at least one chemically reactive function or a chemical group (such as a fluorescent group) which is not present in a natural oligonucleotide and which results
35 from the incorporation of a modified nucleotide or of a non-nucleotide unit carrying this chemically reactive

function or this chemical group. This chemically reactive function makes it possible, inter alia, to perform the synthesis of conjugates of oligonucleotides or of modified oligonucleotides. The terms "chemically reactive function", "modified nucleotide", "non-nucleotide unit" and "oligonucleotide conjugates" are understood to be in the sense described, for example, in the review by J. Goodchild [Conjugates of oligonucleotides and modified oligonucleotides: A review of their synthesis and properties. *Bioconjugate chemistry*, (1990) 1(3), 77-99]. The term "natural oligonucleotide" denotes a polynucleotide formed by the series of nucleotide units existing in nucleic acids [Abbreviations and symbols for the description of conformations of polynucleotide chains. *Eur. J. Biochem.* (1983) 131, 9-15].

The following examples illustrate the invention in a nonlimiting manner.

The following abbreviations are used:

DTT: dithiothreitol
DSS: N-(disuccinimidyl) suberate
GST: glutathione S-transferase
MOPS: 3-[N-morpholino]propanesulfonic acid
SPDP: N-succinimidyl-3-(2-pyridyldithio)propionate
SSMCC: 3-sulfo-N-hydroxysuccinimide ester of 4-(N-maleimidomethyl)cyclohexane-1-carboxylic acid
TEAB: triethylammonium bicarbonate
TEA Ac: triethylammonium acetate containing 10% of acetonitrile.

I/ SYNTHESIS OF CONJUGATES

Example 1: Synthesis of the compounds CY5-T15-hexylamine, CY5-T10-hexylamine and CY5-T5-hexylamine

5

This example describes the synthesis of an oligonucleotide of sequence T₅, T₁₀ or T₁₅ functionalized at its 5' end with a cyanin molecule such as CY5 which is nonsulfonated, and at its 3' end with an arm carrying an
10 amine group which can be used to label a biological molecule of interest. The general structure of the compound CY15-T10 hexylamine for example can be symbolized by ^{5'}(CY5-TTT TTT TTT T-hexylamine)₃.

15 The term "hexylamine" denotes an arm composed of 6 carbon atoms, possibly substituted and carrying an amine function.

In the present example, a 2-hydroxymethyl-6-aminohexanol
20 arm is linked via a phosphate bridge formed between the hydroxyl in the 3' position of the nucleotide located at the 3' end and the 2-hydroxymethyl of the arm.

A solid support of the CPG (controlled pore glass) type
25 conventionally used for synthesizing oligonucleotides is used. Such a support is referred to as "functionalized" since grafted onto the CPG is a chemical structure carrying a protected amine function capable, after final deprotection of the oligonucleotide, of releasing an
30 aliphatic primary amine function.

A commercially available phosphoramidite derivative of thymidine is used for synthesizing the sequence T₁₅.

35 A commercially available phosphoramidite derivative of a nonsulfonated cyanin, which makes it possible to directly

introduce the fluorescent marker such as cyanin (CY5) in the 5' position of the oligonucleotide, is used.

5 The synthesis is carried out using an automatic DNA synthesizer (Applied Biosystems type 392) according to the manufacturer's protocol. The column containing the solid support (CPG) grafted (1 μ mol) with a 2-O-dimethoxytrityl-6-fluorenylmethoxycarbonylamino-1-succinoyl-long chain alkylamino-CPG derivative is placed
10 on the synthesizer, the sequence T₁₅ is synthesized by performing fifteen cycles of synthesis using the phosphoramidite derivative of thymidine, and then a coupling cycle is carried out using the phosphoramidite derivative of the nonsulfonated cyanin (CY5).

15 At the end of this synthesis, the column is subjected to ammoniacal treatment (approximately 2 ml of 28% aqueous ammonia), making it possible to cleave the bond between the oligonucleotide and the CPG support, according to the
20 manufacturer's protocol. The flask for collecting the released oligonucleotide is sealed, kept at 50-55°C for 2 h, and then brought back to ambient temperature. The content of the flask (2 ml) is then transferred into a 5 ml polypropylene tube and then evaporated to dryness
25 under vacuum using a speed-vac. The residue is then taken up with 500 μ l of 10 mM TEAB. The solution obtained contains the "crude" oligonucleotide and predominantly the desired compound ^{5'}(CY5-(T)₁₅-2-oxymethyl-6-amino-
30 hexanol)₃.

The compound ^{5'}(CY5-(T)₁₅-2-oxymethyl-6-amino-
35 hexanol)₃ is obtained after HPLC purification on a LiChrospher RP-18^e 250-10 (10 μ) column (Merck) using a gradient of acetonitrile in aqueous TEAAc (buffer A: 5% acetonitrile in 25 mM TEAAc, buffer B: 50% acetonitrile in 25 mM TEAAc; flow rate 5 ml/min, linear gradient of 10% B to

20% B in 20 min and a linear gradient of 20% to 100% of B in 10 min. The sequences which are incomplete and which do not comprise CY5 are eluted around 20 min, the fractions containing the desired sequence ^{5'}(CY5-(T)₁₅-2-oxymethyl-6-aminohexanol)₃ are collected around 28 min (these fractions are blue in color due to the presence of the CY5 group).

The fractions containing the desired sequence are pooled, and evaporated to dryness using a speed-vac, the residue being taken up with pure water. A UV/visible spectrum (210 nm to 750 nm) effected on a dilution of this solution makes it possible to determine the concentration of the oligonucleotide by its absorbance at 260 nm and to characterize the presence of the cyanin group (CY5) by its absorbance at 650 nm.

The compounds CY5-T10-hexylamine and CY5-T5-hexylamine are synthesized in the same way, by varying the number of cycles of synthesis in the automatic synthesizer.

Example 2: Activation of the compounds CY5-T15-hexylamine, CY5-T10-hexylamine and CY5-T5-hexylamine in the 3' position with SSMCC

The compound CY5-T15-hexylamine obtained in example 1 is dissolved in a 200 mM PO₄ buffer and the pH is adjusted to 8.

250 equivalents of SSMCC are added to the solution obtained. The reaction mixture is incubated for 30 min at ambient temperature, with stirring.

The compound CY5-T15-hexylamine activated with SSMCC, hereinafter referred to as CY5-T15-maleimide (hereinafter named CY5-T15 (mal)), is purified on an HR 10/30 G25 (SF)

column in 10 mM PO₄ buffer containing 2 mM EDTA, pH 7.
The purification is carried out at 60 ml/h.

The solution of CY5-T15-maleimide is concentrated by
5 evaporation in the speed-vac.

The same procedure is carried out for CY5-T10-hexylamine.

For the activation of the compound CY5-T5-hexylamine,
10 only 150 equivalents of SSMCC are used. An additional
purification step on an HR 10/30 column is necessary.

Example 3: Activation of CY5-T15-hexylamine in the 3'
position with DSS (N-(disuccinimidyl suberate)

15

The compound CY5-T15-hexylamine obtained in example 1 is
taken up in 10 µl of 100 mM MOPS buffer, pH 7.6, and 5 µl
of acetonitrile.

20 150 equivalents of DSS (35 mg/ml in DMF) are added to the
solution obtained.

The reaction mixture is incubated overnight at 4°C with
stirring.

25 The compound CY5-T15-hexylamine activated with DSS,
hereinafter referred to as CY5-T15-NHS, is purified on a
NAP 5 G25 (SF) column in 5 mM MOPS buffer, pH 6.5.

30 The compound CY5-T15-NHS is then concentrated by several
precipitations with butanol and centrifugations. The
pellet is taken up in water.

The same procedure is carried out for the activation of
the compounds CY5-T10-hexylamine and CY5-T5-hexylamine.

35

Example 4: Preparation of the conjugate CY5-(T)_n-hexyl-amine activated with SSMCC (example 2) - anti-GST antibody (GSS11)

5 **GSS11-T15-CY5 batch 02B (mal):**

The antibody GSS11 (CIS bio international, France) in 0.1 M carbonate buffer, pH 9, is activated by adding 8 equivalents of SPDP for 30 min at ambient temperature,
10 with stirring, and then adding a final concentration of 20 mM of DTT for 15 min at ambient temperature, without stirring.

It is purified on an HR 10/10 G25 (SF) column in 0.1 M
15 PO₄ buffer, pH 7.

The antibody thus activated is mixed with the oligonucleotide CY5-T15-maleimide obtained in example 2, with an initial CY5-T15-maleimide/antibody GSS11 molar
20 ratio of 7.6. The incubation is from 18 to 20 h at +4°C.

The concentration of the antibody during the coupling is 0.5 mg/ml.

25 **GSS11-T15-CY5 batch 03 (mal):**

The antibody GSS11 is activated only by adding DTT at a concentration of 5 mM.

The same procedure as for batch 02B above is then carried
30 out, with an initial CY5-T15-maleimide/antibody GSS11 molar ratio of 10.8.

The concentration of the antibody during the coupling is, in this case, 0.68 mg/ml.

GSS11-T5-CY5 batch 01 (mal):

The same procedure as for batch 02B above is carried out, with an initial CY5-T5-maleimide/antibody GSS11 molar ratio of 8.

The concentration of the antibody during the coupling is 0.9 mg/ml.

10 GSS11-T10-CY5 batch 01 (mal):

The same procedure as for the GSS11-T5-CY5 batch 01 (mal) is carried out.

15 The coupling products are purified on an HR 10/30 Superdex 200 column at 60 ml/h.

The elution buffer is a 0.1 M phosphate buffer, pH 7.

20 The purifications are followed using a diode array detector (followed at various wavelengths).

Example 5: Preparation of the conjugate CY5-(T)_n-hexylamine activated with DSS (example 3) - anti-GST antibody (GSS11)

GSS11-T15-CY5 batch 01 (NHS):

30 The antibody GSS11 in 0.1 M carbonate buffer, pH 9, is mixed with the CY5-T15-NHS solution obtained in example 3. The initial CY5-T15-NHS/antibody molar ratio is 8. The concentration of the antibody during the coupling is 3.3 mg/ml. The incubation is for 30 min at ambient temperature without stirring.

35

GSS11-T15-CY5 batch 02 (NHS):

The protocol is the same as for GSS11-T15-CY5 batch 01 NHS, with an initial CY5-T15-NHS/antibody molar ratio of
5 12. The incubation is for 2 h 30 at ambient temperature, but with stirring.

The coupling products are purified on an HR 10/30 Superdex 200 column at 60 ml/h. The elution buffer is a
10 0.1 M phosphate buffer, pH 7.

The purifications are followed using a diode array detector (followed at various wavelengths).

15 **Example 6: Synthesis of the conjugate CY5sulfo-GSS11**

These conjugates, which do not comprise any oligonucleotide, serve as reference compounds to show the advantages of the conjugates according to the invention.

20

A sulfonated cyanin is used here, and not a nonsulfonated cyanin as in the previous examples, since the quantum yield of the latter is virtually zero if it is not coupled to an oligonucleotide.

25

GSS11-CY5sulfo batch M5:

Sulfonated cyanin (CY5sulfo) mono NHS is added to antibody GSS11 in 0.1 M carbonate buffer, pH 9.

30

The initial CY5sulfo/antibody molar ratio is 4.

The incubation is for 1 h at ambient temperature with stirring, the concentration of the antibody during the
35 coupling is 5.2 mg/ml.

The purification is carried out on an HR10/10 G25 (SF) column in 0.1 M phosphate buffer, pH 7.

GSS11-CY5sulfo batch M6:

5

The protocol is the same as for batch M5 above, but with an initial CY5sulfo/antibody molar ratio of 10.

GSS11-CY5sulfo batch M7:

10

The protocol is the same as for batch M5 above, but with an initial CY5sulfo/antibody molar ratio of 20.

Example 7: Preparation of the conjugate CY5-A15-cAMP

15

The compound Cy5-A15-hexylamine is obtained according to a procedure similar to that described for the Cy5-T15-hexylamine in example 1, using a phosphoramidite derivative of adenosine in place of the phosphoramidite derivative of thymidine.

20

The synthesis of the N-hydroxysuccinimide ester of 2'-monosuccinyladenosine 3',5'-cyclic monophosphate (cAMP-succ-NHS) is carried out as follows:

25

2 mg (3.9 μ mol) of 2'-monosuccinyladenosine 3',5'-cyclic monophosphate (sodium salt, Sigma # M9631) are suspended in 15 μ l of a solution of N-hydroxysuccinimide (13.3 mg/ml in anhydrous DMSO), in an eppendorff tube.

30

35 μ l of a solution of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (200 mg/ml in anhydrous DMSO) are then added to the reaction volume and then placed on a shaker at ambient temperature.

35

After reaction for 90 minutes, the reaction mixture is diluted with DMSO qs 160 μ l, so as to obtain a solution of [cAMP-succ-NHS] = 23 nmol/ μ l, which is used for the synthesis of conjugates.

5

The cAMP-succ-NHS taken up in H₂O is then mixed with a solution of CY5-A15-hexylamine in 10 mM TEAB buffer, pH = 8.5. During the coupling reaction, the molar ratio is 20 cAMP-NHS per CY5-A15-hexylamine. The cAMP
10 concentration during the coupling is 5 μ mol/ml. The incubation is for 2 h 30 at ambient temperature with agitation.

The purification is carried out using desalification on a
15 Nap5 (G25) column in 0.1 M PO₄ buffer, pH 7.

II - PHOTOPHYSICAL PROPERTIES

Example 8: Comparison of the quantum yields and of the OD
20 max/OD 604 nm ratio

The quantum yield reflects the efficiency of the fluorescent entity in releasing the energy which it receives: the higher it is, the more efficient the
25 fluorescent entity. These quantum yields are measured using a fluorimeter. The excitation is carried out at 600 nm, the fluorescence is measured at 615 to 750 nm. The area of the fluorescence spectra obtained is calculated and used to determine the quantum yields.

30

The final molar ratio (FMR) expresses the number of fluorophores covalently coupled to the protein.

The absorption spectra of the fluorescent entities make
35 it possible to calculate the OD max/OD 604 nm ratio. This ratio reflects a possible phenomenon of aggregation of

the fluorophores, for example CY5, at the surface of the labeled proteins, for example the antibody GSS11 or streptavidin.

- 5 The quantum yield and the OD max/OD 604 nm ratio are determined for various compounds synthesized according to the preceding examples.

8.1/ The results given in table 1 below concern the
10 reference conjugates (comprising no oligonucleotide) prepared in example 6 above.

Table 1

	Final molar ratio (FMR)	OD _{max} / OD _{604 nm}	Quantum yield
CY5sulfo-mono NHS	-----	3.16	19%
GSS11-CY5sulfo batch M5	2.20	2.18	11%
GSS11-CY5sulfo batch M6	3.80	1.64	4%
GSS11-CY5sulfo batch M7	6.00	1.24	1%

15

These results show that the coupling of the sulfonated cyanin CY5sulfo with an antibody GSS11 leads to a decrease in the quantum yield, associated in particular with a phenomenon of aggregation of CY5 at the surface of
20 the antibody, as shown by the decrease in the ODmax/OD 604 nm ratio.

The more the number of CY5 molecules per antibody molecule (FMR) increases, the greater this decrease.

25

8.2/ The results given in table 2 below concern the conjugates prepared in examples 4 and 5. The compounds CY5-T10-hexylamine and CY5-T15-hexylamine used as references are prepared as described in example 1.

30

Table 2

	Final molar ratio (FMR)	OD _{max} / OD _{604 nm}	Quantum yield
CY5-T15-hexylamine	-----	2.79	25%
GSS11-T15-CY5 batch 02B (mal)	4.30	2.59	19%
GSS11-T15-CY5 batch 03 (mal)	1.40	2.59	18%
GSS11-T15-CY5 batch 01 (NHS)	2.00	2.67	20%
GSS11-T15-CY5 batch 02 (NHS)	5.10	2.63	20%
CY5-T10-hexylamine	-----	2.78	26%
GSS11-T10-CY5 batch 01 (mal)	4.60	2.38	14%

These results show that, when coupling the antibody GSS11
5 with fluorescent entities according to the invention, a
very small decrease in the quantum yield and in the OD
max/OD 604 nm ratio is observed, in comparison to the
reference conjugate, for increasing FMRs. This clearly
shows that the fluorescent entities according to the
10 invention exhibit completely unexpected properties in
terms of decreasing the aggregation at the surface of the
labeled protein (here the antibody GSS11).

In addition, the result obtained is independent of the
15 method of activation of the compound CY5-T15-hexylamine
(DSS or SSMCC).

8.3/ The results given in table 3 below concern the
product prepared in example 7. The compound Cy5-A15-
20 hexylamine used as a reference is prepared according to
the procedure described for Cy5-T15-hexylamine in example
1.

Table 3

	OD _{max} /OD ₆₀₄	Quantum yield
CY5-A15-NH2	2.7	16%
CY5-A15-cAMP	2.65	19.5%

The results above show that the quantum yield of fluorescence of the fluorescent entities is slightly increased after coupling to the cyclic AMP. The conjugate thus created exhibits no sign of aggregation and thus has unexpected fluorescence properties.

10 **Example 9: Fluorescence spectrum at constant antibody concentration**

In order to compare the level of fluorescence obtained with the various conjugates, fluorescence emission spectra (intensity of fluorescence as a function of the emission wavelength) were produced with a constant concentration of antibody (50 nM). These spectra are produced on an LS50B spectrofluorimeter (PerkinElmer), with the settings as follows: excitation wavelength 20 600 nm, emission wavelength 615 to 750 nm, scanning rate 480 nm/min. Based on these spectra, it is possible to determine the intensity of fluorescence of the conjugate by calculating the area of the spectrum obtained by integration.

25 Figure 1 gives the relationship between the FMR of the conjugates GSS11-CY5sulfo and GSS11-T15-CY5 and their intensity of fluorescence.

30 The following symbols are used:

- represents the conjugate GSS11-CY5 sulfo
- represents the conjugate GSS11-T15-CY5

The graph in Figure 1 shows that, in the case of the conjugates GSS11-CY5sulfo, the increase in the FMR of the conjugate leads to a decrease in its overall fluorescence, due to the extreme aggregation of the CY5 at the surface of the antibody. On the other hand, in the case of the conjugates GSS11-T15-CY5, the absence of aggregation of the CY5 makes it possible to maintain a high quantum yield and, consequently, to obtain an overall fluorescence which is virtually proportional to the number of CY5 per antibody.

III - USE OF THE FLUORESCENT ENTITIES ACCORDING TO THE INVENTION IN ASSAYS OF THE FRET ("Fluorescence Resonance Energy Transfer") TYPE

Example 10:

The fluorescent entities according to the invention can be used in systems of the FRET type well known to those skilled in the art.

In the present example, biotinylated Glutathione S-transferase (GST-biotin) is detected by measuring the fluorescence emitted by an acceptor compound, resulting from an energy transfer between a donor compound (conjugate europium cryptate-streptavidin (K(Eu)-Sa)) and an acceptor containing a fluorescent entity according to the invention (conjugate GSS11-oligonucleotide-CY5).

Assay protocol:

The assay is carried out using a fluorimeter (Discovery, Packard), the excitation wavelength of which is 337 nm. The fluorescence is measured at 665 and 620 nm.

Assay buffer: 50 mM Hepes, pH 7, 0.1% BSA, 400 mM KF.

Reagents: GST-biotin, 20 nM solution

Donor: Sa-K(EU) NHS (CIS bio international)

5 Acceptor: GSS11-XL665 (CIS bio international) used as reference

GSS11-T15-CY5 batch 02B mal (ex. 4)	FMR 4.3
GSS11-T15-CY5 batch 03mal (ex. 4)	FMR 1.4
GSS11-T15-CY5 batch 01NHS (ex. 5)	FMR 2
GSS11-T15-CY5 batch 02NHS (ex. 5)	FMR 5.1
GSS11-T10-CY5 batch 01mal (ex. 4)	FMR 4.6
GSS11-T15-CY5 batch 01mal (ex. 4)	FMR 8.7

The following mixture is incubated for 20 h at ambient temperature:

- 10 - 50 µl of GST-biotin at 0 - 0.31 - 0.62 - 1.25 - 2.5 or 5 nM final concentration
- 100 µl of acceptor at 2.5 nM final concentration
- 50 µl of donor at 1 nM final concentration

15 Figure 2 represents the evolution of the signal (% delta F) as a function of the evolution of the concentration of the GST-biotin (GST-BIOT in nM final concentration).

The following symbols are used:

- 20 —○— GSS11-XL-665
- ◆— represents the conjugate GSS11-T15-CY5 batch 02 NHS
- — represents the conjugate GSS11-T15-CY5 batch 01 NHS
- 25 —△— represents the conjugate GSS11-T15-CY5 batch 02B mal
- x— represents the conjugate GSS11-T15-CY5 batch 03 mal
- — represents the conjugate GSS11-T10-CY5 batch 01 mal
- 30

The graph in Figure 2 shows the advantage of the fluorescent entities according to the invention as fluorescent labels in an assay of the FRET type. Specifically, in all cases, the signal observed using the
5 fluorescent entities according to the invention is greater than or equal to that obtained with the reference acceptor compound (XL).

Example 11: Lifetime and efficiency of transfer of the
10 **conjugates**

The lifetimes and the efficiency of transfer obtained in the assay of the FRET type such as that of the preceding example were calculated. The conjugates tested were
15 prepared as described in examples 4 and 5, the conjugate GSS11-XL665 serving as a reference.

The results are given in table 4 below.

Table 4

	Sa-K batch 14 (NHS generic product)		
	Lifetime of the FRET in ms (τ_{FRET})	Lifetime of the free K in ms (τ_{cryptate})	Efficiency of transfer in % $1 - (\tau_{\text{FRET}}/\tau_{\text{cryptate}})$
GSS11-T10-CY5 batch 01mal	0.16	0.969	83%
GSS11-T5-CY5 batch 01mal	0.21	1.030	80%
GSS11-T15-CY5 batch 03mal	0.21	1.054	79%
GSS11-T5-CY5 batch 02mal	0.16	1.001	83%
GSS11-T15-CY5 batch 01NHS	0.17	0.997	83%
GSS11-T15-CY5 batch 02NHS	0.13	1.038	87%
GSS11-XL665	0.27	1.033	73%

5 The results show that the efficiency of energy transfer between the donor compound and the conjugates according to the invention used as acceptors is significantly higher for the conjugates according to the invention than for the conjugate XL665 (control).

10 Example 12:

The fluorescent entities can be used in FRET competition systems well known to those skilled in the art.

15 In the present example, the presence of cyclic AMP (cAMP) in a sample is detected by observing the inhibition of the fluorescence energy transfer occurring between a

donor compound (conjugate europium cryptate-anti-cAMP monoclonal antibody) and an acceptor compound containing a fluorescent entity (conjugate Cy5-A15-cAMP).

5 Assay protocol:

The assay is carried out using a fluorimeter (Rubystar, BMG), the excitation wavelength of which is 337 nm. The fluorescence is measured at 665 nm and 620 nm.

10

Assay buffer: 0.1 M phosphate, pH = 7, 0.1% BSA, 400 mM KF.

Reagents: cAMP, solution at 280 nM

15 Donor: conjugate Europium cryptate-anti-cAMP monoclonal antibody (anti-cAMP-K) (Cis bio international)
Acceptor:

conjugate Cy5-A15-cAMP (Cis bio international),
conjugate cAMP-XL665 used as a reference.

20

The following mixture is incubated for 20 h at ambient temperature:

- 25 µl buffer
- 25 µl cAMP at 0-0.07-0.27-1.09-4.37-17.5 or 70 nM
- 25 final concentration
- 25 µl cAMP-XL665 or Cy5-A15-cAMP
- 25 µl anti-cAMP-K

30 Figure 3 represents the inhibition of FRET signal (DF/DF max) obtained at two incubation times (1 h and 20 h) in the presence of increasing amounts of cAMP.

The following symbols are used:

- --- cAMP-XL665 incubation time 1H
- 35 ---o--- cAMP-XL665 incubation time 20H
- x--- Cy5-T15-cAMP incubation time 1H

—▽— Cy5-T15-cAMP incubation time 20H

The graph in Figure 3 shows the advantage of the fluorescent entities according to the invention as
5 fluorescent labels in an assay of the FRET competition type. The sensitivity of the test is improved by using the fluorescent entities according to the invention, whatever the incubation time of the experiment. In
10 addition, the loss of sensitivity observed with the reference acceptor (XL665) between 1 h and 20 h of incubation disappears when the fluorescent entities according to the invention are used.